From seed to supermassive

growing high redshift black holes in cosmological simulations

Ricarda Beckmann

working with Julien Devriendt and Adrianne Slyz University of Oxford




Feedback from massive black holes is responsible for quenched galaxies in the local universe


Effect of AGN feedback

Data from the HORIZON-AGN & HORIZON-noAGN simulations: Dubois et al 2014, Welker et al 2015, Volonteri et al 2016, Kaviraj et al 2016, Beckmann et al (in prep)

At z=6, massive galaxies are already quenched



Black holes growth precedes galaxy growth



Black holes in cosmological simulation

In HORIZON (Dubois2014,Welker2015,...):

- Seed the BHs at 1E5 Msun
- 1kpc resolution
- Grow using Eddington limited Bondi accretion

=> Similar picture in other simulations such as Illustris (Vogelsberger2014,Sijaki2015), Eagle (Schaye2015) and massive black II (Khandai2014)


Where do these black holes physically comes from?

SEEDING MODELS


- **Direct Collapse:** 1E5 Msun seeds, needs metal free gas and radiating neighbours (Begelman2006,Latif2013,Habouzit2016)
- Runaway cluster collapse: 1E3-1E4 Msun seeds (Omukai2008, Devecchi2009,Katz2015)
- Stellar mass seeds: 1-100 Msun, (Haiman2000, Freyer2001, Alvarez2009, Heger2003)


Bridging the gap from seed to supermassive

- stellar mass seed (260 MSun)
- a cosmological context (1Gpc box with 5 Mpc zoom region)
- high resolution ($\Delta x=0.01$ pc)
- run to redshift 6
- AMR hydro (RAMSES)




Zoom-within-Zoom






Gradual triggering of levels




Density




RESULTS


A test case: an isolated disc




A test case: a sink in an isolated disc




A test case: small scale features



The cosmological simulation



The cosmological simulation



Thank you